

Reduction of acrylamide formation in cereal-based food  
processing

5           The present invention relates to improvements in  
and relating to cooked food, in particular cereal-based  
foods which are fried, grilled, baked or roasted.

10           In a publication by the Swedish National Food  
Administration (see [www.slv.se/engdefault.asp](http://www.slv.se/engdefault.asp)) it was  
reported that many cooked foods, in particular fried,  
grilled or baked foods, had surprisingly been found to  
contain high levels of the toxic contaminant acrylamide.  
No suggestion was made as to how the acrylamide context  
of such foods could be reduced.

15           A further report of acrylamide production in food  
cooking occurred in Tareke et al., J. Agric. Food Chem  
50: 4998-5006 (2002).

20           We have now surprisingly found that the acrylamide  
content of cooked cereal-based foods can be reduced by  
treatment of the food or the cereal base therefor prior  
to cooking with lactic acid generating microorganisms  
and/or with acid.

25           Thus viewed from one aspect the invention provides  
the use of a lactic acid producing microorganism for the  
treatment of a cereal-based food material or the cereal  
base therefor to reduce acrylamide production in  
subsequent cooking thereof.

30           Lactic acid producing microorganisms are well known  
and examples include lactic acid bacteria such as  
Bifidobacterium sp., Brevibacterium sp., Lactobacillus  
sp., Lactococcus sp., Leuconostoc sp., Micrococcus sp.,  
Oenococcus sp., Pediococcus sp., and Streptococcus sp.  
Lactobacilli are especially preferred for use according  
to the invention, in particular Lactobacillus plantarum  
35           strains NCDO 1752 and NCDO 1193 (available from the  
National Collection of Food Bacteria) and Lactobacillus  
NCIMB 40450. Other strains of lactobacillus which

- 2 -

generate lactic acid and are safe for use in foodstuff treatment have been described widely in the scientific literature.

5 The treatment with a lactic acid producing microorganism according to the invention preferably involves incubation in an aqueous medium for up to 7 days, e.g. 30 minutes to 24 hours, especially 1 to 6 hours. Incubation is preferably at 4 to 45°C, e.g. 25 to 35°C, i.e. as is conventional for such  
10 microorganisms.

Typically such treatment may involve homofermentative lactic acid bacteria incubation in an aqueous medium.

15 Viewed from a further aspect the invention provides the use of a physiologically acceptable acid for the treatment of a cereal-based food material or the cereal base therefor to reduce acrylamide production in subsequent cooking thereof.

20 The physiologically tolerable acid used according to the invention may be any acid acceptable for use in foodstuffs, e.g. organic acids, such as citric, malic, acetic, maleic, tartaric, succinic and lactic acids or inorganic acids such as hydrochloric, sulphuric and phosphoric acids and sulphur dioxide. The use of citric  
25 and hydrochloric acids is especially preferred, as is the use of lactic acid and/or of phosphoric acid. The use of hydrochloric acid and of lactic acid is particularly preferred. The acid is preferably used in a quantity and strength sufficient to reduce the surface  
30 pH of the food material treated to 1 to 5.5, preferably 3 to 5, especially about 4. Following acid treatment, the food material is preferably stored for up to 7 days (e.g. 30 minutes to 24 hours, especially 1 to 6 hours before cooking or freezing.

35 In this process, the acid is preferably used in the form of a buffer solution.

Following treatment with the acid and/or the lactic

- 3 -

acid producing microorganism, the food material may be cooked using cooking techniques that expose the material to temperatures above 150°C, e.g. by baking, grilling, roasting or frying.

5        Before such high temperature cooking, it is desirable to rinse the treated food material with water.

         The cooking may be a single stage operation. However it may instead be one stage of a multi stage (e.g. two stage) cooking procedure. Thus the technique  
10        of the invention is especially applicable to food materials which are treated according to the invention, partially cooked, transported and/or stored, then cooked again.

         The invention is particularly applicable to  
15        products made from cereal (e.g. rice, barley, wheat, rye, oat, maize, etc.) flours, granulates or fragments, in particular breads (especially crisp-breads, biscuits, wafers, cookies and crackers), cakes, snacks (e.g. crisps (in American-English chips), pretzels, and the  
20        like), and breakfast cereals (e.g. "cornflakes" and the like).

         Thus in a further aspect the invention provides a process for the production of a food product which comprises fermenting a granulated or crushed  
25        carbohydrate-containing cereal material with a lactic acid producing microorganism, optionally formulating the fermented material into a shaped product (e.g. by extrusion, rolling or moulding a paste or dough), and cooking to produce said food product.

30        In place of fermentation, acid treatment as described above may be used: however this is less preferable.

         The granulated carbohydrate-containing cereal material, e.g. a cereal flour, may be mixed with un-  
35        treated granulated carbohydrate-containing cereal material before cooking. Desirably the treated:untreated weight ratio is from 25:75 to 100:0,

- 4 -

especially 50:50 to 95:5.

The term granulate or granulated as used herein may where the context permits include fine to coarse particulates, e.g. flours, granules, grits, fragments, etc. Preferably however the granulates will be 2mm or smaller in maximum dimension.

The food products thus produced are desirably packed into sealed, preferably sterilized containers, e.g. cartons, plastic or foil bags, bottles, etc. Particularly preferably such containers are moisture-proof.

Where appropriate, the food products produced according to the invention may be further processed, e.g. by drying, freezing, cutting, decorating, etc. Such processing steps, which are often conventional for the particular food product, form further optional steps in the processes of the invention.

French-fries may also be made from extruded or moulded carbohydrate-containing pastes produced using powdered or granulated cereal (e.g. rice) optionally together with powdered or granulated potato.

The invention is applicable to breads, biscuits (known in America as cookies), and in particular crisp breads. In this aspect of the invention, the treatment according to the invention may be effected using lactic acid bacteria in the production of the dough and/or by acid treatment (e.g. treatment with sulphur dioxide or hydrogen chloride) of the flour.

Besides the fermentation and/or acid treatment according to the invention, the food products of the invention may be prepared by conventional methods, optionally involving rinsing and/or drying after the treatment. Thus such food products may optionally contain further components, such as conventional foodstuff components or additives, e.g. salt, sugars, flavours, fruit, fruit extracts, nuts, eggs, milk, flour, bread, breadcrumbs, stabilizers, colours,

- 5 -

buffers, acidulants, yeast, bicarbonate, etc.

The invention will now be illustrated further with reference to the following non-limiting Example.

5     Example 1

Crisp bread

Ingredients

Finely ground whole rye flour was obtained from Cerealia Mills, Oslo, Norway.

10

Crisp bread recipe

The crisp bread dough was based on a standard recipe made from 1150g rye flour, 1000g water, 20g NaCl. The ingredients were mixed and the dough was rolled out to proper thickness and baked in a stone oven at 240°C.

15

Acid equivalent

Total amount of acid in the flour-water mixtures was determined as Acid equivalents, S°, by titration. To 10g flour-water mixture, 90g of distilled water is added. During stirring, 0.1 N NaOH is added until a stable pH of 8.5 is reached. Acid equivalents is expressed as the amount of 0.1 N NaOH consumed, in ml.

20

25     Pre-treatments

Fermentation

The rye flour used in the crisp bread dough was fermented using a lactic acid bacteria (e.g. NCIMB 40450 or others described herein) according to the following description: 1000g rye flour and 1000g tap water at 30°C was mixed and bacteria (10<sup>6</sup> bacteria/g flour) was added. The bacteria had previously been cultured in MRS growth medium, harvested in the exponential phase, centrifuged and dispersed in water, prior to being added to the flour. Flour fermentation was performed at 30°C in a proofing cabinet at 70% relative humidity (RH) for about 18 hours.

30

35

- 6 -

*Soaking in lactic acid*

The rye flour used in the crisp bread was soaked in lactic acid according to the following description:  
1000g flour was mixed with 1000g 0.15 M lactic acid at  
30°C.

*Soaking in acidic phosphate buffer*

The rye flour used in the rye crisp bread was soaked in phosphate buffer according to the following description:  
1000g flour was mixed with 1000g 0.1 M phosphate buffer, pH 4.0, 30°C.

Doughs with the following composition were produced from the pre-treated or non-pre-treated flours:

1. No pre-treatment

345g rye flour + 300g water + 7g NaCl was mixed for 4 min in a Hobardt mixer equipped with a dough hook.

2. Fermentation - lactic acid bacteria added

450g fermented flour/water mixture (bacteria added) + 150g untreated rye flour + 7g NaCl was mixed for 4 min in a Hobardt mixer equipped with a dough hook.

3. Lactic acid

450g flour/water mixture in lactic acid + 150g untreated rye flour + 7g NaCl was mixed for 4 min in a Hobardt mixture equipped with a dough hook.

4. Phosphate buffer

450g flour/water mixture in phosphate buffer + 150g untreated rye flour + 7g NaCl was mixed for 4 min in a Hobardt mixer equipped with a dough hook.

The proportion of pre-treated flour in doughs 2 to 4 was thus 60%.

- 7 -

About 100g of dough was rolled to 0.5mm thickness and baked at 240°C for 10 min.

Analysis of acrylamide was effected by Norsk Matanalyse AS, Oslo, Norway.

### Results

The results are presented in Table 1 below.

10	Pre-treatment of rye flour	pH after pre- treatment	Acid equivalent		Acrylamide	
			ml 0.1N NaOH/10g	µg/kg product	% reduction*	(ppb)
15	No-pretreatment	6.00	4.0	3500	-	
	Fermentation - lactic acid bacteria added	3.70	18	890	75	
	Soaked in lactic acid	4.05	20	1000	71	
20	Soaked in phosphate buffer	4.74	11	1400	60	

\*Reduction relative to crisp bread with no pre-treatment.

The pH in the rye flour/water mixture with bacteria added decreased during the pre-treatment. This shows that fermentation had occurred with corresponding acid production. The pH of the mixture with lactic acid and phosphate buffer added was as expected. At the applied conditions, adding lactic acid was effective in lowering pH and was thus similar to the controlled fermentation. The acid equivalent values reflected the changes seen in pH, with higher levels corresponding to the most extensive changes in pH.

Acrylamide levels in crisp bread reflected the various

- 8 -

- pre-treatments tested. With no pre-treatment, the acrylamide level was high. The use of 60% rye flour fermented by the adding of lactic acid bacteria caused a 75% reduction in acrylamide compared with crisp bread produced without pre-treatment. The reduction was 71% when 60% of the flour was soaked in lactic acid as a pre-treatment. With phosphate buffer, the reduction in acrylamide was 60%.
- 10 It is likely that the pre-treatment can be optimised in several ways. First, the present experiments were performed with pre-treatment of only 60% of the total flour in the crisp bread. Doughs based on 100%-fermented flour may be used. Alternatively, the
- 15 fermented flour may be fully or partly dried prior to application in the crisp bread dough, thus making it possible to adjust the viscosity of the dough to a level appropriate for rolling. Combinations of fermentation and acids may be effective in further lowering
- 20 acrylamide levels, as well as sole adding of lactic or other acids to higher levels, and/or lower pH values.